



HAL
open science

Atmospheric ammonia in-situ long-term monitoring: review worldwide strategies and recommendations for implementation

Aude Bourin, Pablo Espina-Martin, Anna Font, Sabine Crunaire, Marsailidh M Twigg, Christine F Braban, Stéphane Sauvage

► To cite this version:

Aude Bourin, Pablo Espina-Martin, Anna Font, Sabine Crunaire, Marsailidh M Twigg, et al.. Atmospheric ammonia in-situ long-term monitoring: review worldwide strategies and recommendations for implementation. EGU General Assembly 2023, Apr 2023, Vienne, France. 2023, 10.5194/egusphere-egu23-2847 . hal-04199511

HAL Id: hal-04199511


<https://imt-nord-europe.hal.science/hal-04199511v1>

Submitted on 8 Sep 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Ammonia (NH₃): Atmospheric role




- Major alkaline gas in the atmosphere: mainly agricultural sources.
- Forms secondary salts, mostly on the fine particle fraction (PM_{2.5}).
- Dry and wet deposition: causes damage on ecosystems, eutrophication and acidification issues.
- Recent studies suggests it enhances new particle formation.

Reductions in NH₃ emissions and ambient concentrations are necessary to improve air quality (AQ) and recover sensitive ecosystems¹. Worldwide there are relatively few long-term national and continental NH₃ concentration monitoring networks. There is not yet information from South America, Southwestern Asia, large parts of Africa and Oceania. The aim of this study is to review current networks and provide recommendations for setting up future NH₃ long-term monitoring.

Long-term monitoring networks: Main objectives pursued

1. Estimate regional NH₃ concentrations and establish national NH₃ baselines.
2. Assess spatial & temporal patterns at short-term and long-term driving NH₃ concentration variability.
3. Estimate impacts caused by NH_x deposition in ecosystems and protected areas.
4. Apportion the origin of the reduced nitrogen (N_{red}) deposition and/or estimate the N_{red} deposition fluxes.
5. Assess the effectiveness of mitigation strategies and inform the grand public on AQ changes.
6. Validate and/or calibrate models (chemical transport, dispersion, emissions...).

Long-term monitoring networks of NH₃ worldwide



	Network	Temporal coverage	N sites	Sampling method (A: active, P: passive)	Temporal resolution	Objectives
USA ²	AMON	2007-Present	106	Radiello tubes (P)	Biweekly	1,3,4,6
Canada ²	NAPS	2003-present	14	Denuder filterpack (A)	3 -6 days	1,2,4,5
Europe-Asia ³	EMEP	1986 - Present	TBD*	Various (A, P)	Various	2,5,6
UK ⁴	UKEAP NAMN AGA-Net EMEP supersites	1998-Present	78	ALPHA sampler (P)	Monthly	1,2,3,6
			28	DELTA (A)	Monthly	
			2	MARGA (A)	1h	
Netherlands ⁴	LML MAN	1993-Present 2005-Present	6	miniDOAS (A)	1h	2
			>300	Gradko sampler (P)	Monthly	1,3,6
Ireland ⁴	NEMN	2017-Present	TBD*	ALPHA sampler (P) DELTA (A)	Monthly	2,3,5
Switzerland ⁴	-	2000-Present	>80	Ferm sampler (P)	2-4 weeks	2,5,6
Belgium ⁴	-	2002-Present	18	Radiello sampler (P) Mini-DOAS (A)	Monthly 1h	1,2,3,6
China ⁵	AMON	2015-Present	53	Analyst sampler (P)	Monthly	2
Asian Southeast ⁶	EANET	1999-Present	44	Denuder filterpack (A)	Weekly, 10 days, Biweekly	2,3,5
Western Africa ⁷	INDAAF	1995-Present	13	INDAAF sampler (P)	Monthly	2,4

TBD*: to be determined, bibliography research not conclusive

Recommendations for future NH₃ monitoring networks

In the near future, the number of NH₃ monitoring networks using FAIR data-archiving is expected to increase to better evaluate AQ objectives and follow the long-term changes in N_{red} deposition. In this poster, we propose some recommendations and approaches used by current long term monitoring networks:

1. Co-locate with established monitoring sites to minimise costs, where possible.
2. Smaller countries may share monitoring sites with surrounding countries to keep running costs low.
3. Tiering monitoring sites by mixing passive monitoring methods with high temporal monitoring methods provides spatial distribution and evidence of the sources and processes driving the reported NH₃ concentrations.
4. The spatial strategy of having a high number of monitoring sites equipped with passive samplers allow to achieve objectives 1,2,3, and 5. Local site operators (LSOs) can help identify local sources of NH₃ and can potentially decrease the operational costs.

5. High temporal resolution active monitoring is advised to identify NH₃ temporal changes drivers and, within ACTRIS, EMEP and WMO-GAW, traceable protocols should be developed and tested and include pre-existing networks using continuous wet methods (MARGA in UKEAP), spectroscopic methods (miniDOAS in LML & Belgium) and other new technologies.
6. Data should be made open access via an online data repository, where possible. Information's such as the network operational conditions and QA/QC protocols are also beneficial.

References & Acknowledgments

1: Jonson, J. E., et al.; Modelling changes in secondary inorganic aerosol formation and nitrogen deposition in Europe from 2005 to 2030, *Atm. Chem. Phys.*, 22, 1311–1331, 2022.
 2: Nair, A. A. and Yu, F.: Quantification of Atmospheric NH₃ concentrations: A Review of Its Measurement and Modeling, *Atmosphere*, 11, 1092, 2020.
 3: Hjelbrekke, A.-G.: EMEP/CCC-Report 1/2022 on Particulate matter, carbonaceous and inorganic compounds, Norwegian Institute for Air Research, 2022.
 4: Franzaring, J. and Köstler, J.: Review of internationally proposed critical levels for ammonia, University of Hohenheim, Stuttgart, 2022.
 5: Pan, Y. et al.: Identifying Ammonia Hotspots in China Using a National Observation Network, *Environ. Sci. Technol.*, 52, 3926–3934, 2018.
 6: Hatakeyama, S., et al.; The Fourth Periodic Report on the State of Acid Deposition in East Asia Part I: Regional Assessment, 2021.
 7: Adon, M., et al.: Long term measurements of sulfur dioxide, nitrogen dioxide, ammonia, nitric acid and ozone in Africa using passive samplers, *Atm. Chem. Phys.*, 10, 7467–7487, 2010.
 This work has been supported by the French Ministry of Environment, France as a part of the activities of French observatory for the measurement and evaluation of long range air pollution in rural areas (MERA). Thanks to Jeroen Staelens (VMM) for the information on the Flanders region NH₃ network, to David Kelleghan (EPA Ireland) for the information related to the Irish NEMN, to Ewa Dabek-Zlotorzynska and Valbona Celso for the information related to the NAPS program in Canada, to Christoph Hüglin, Reto Meier and Eva Seitler for providing insight on the Swiss NH₃ network.